

ON $F_a(K, (K-2))$ -STRUCTURE MANIFOLD

Jai Pratap Singh

B.S.N.V.P.G. College, Lucknow, India Email: jaisinghjs@gmail.com

Kripa Sindhu Prasad

Department of Mathematics, Thakur Ram Multiple Campus, Birgunj, Tribhuvan Univeristy, Nepal Email: kripasindhuchaudhary@gmail.com

Abstract

Yano, Houh and Chen [1] have studied the structure defined by the tenosor field ϕ of type (1,1) satisfying $\phi^4 \pm \phi^2 = 0$. Gadea and Cordero [2] have obtained the integrability conditions of these structures. The purpose of this paper is to define and study $F_a(K, (k-2))$ -structure. Integrability conditions of such a structure have also been studied.

Keywords: Riemannian Manifold, projection operator, invariant submanifold, integrability conditions **AMS Classification:** 53C05, 53C15

References

- [1] 10 K.Yano, C.S. Houh and B.Y. Chen (1972). Structure defined by a tensor field ϕ of type (1,1) satisfying $\phi^4 \pm \phi^2 = 0$ Tensor, NS, vol. 23, pp. 81-87.
- [2] P.M. Gadea and L.A. Cordero (1974). Integrability conditions of a structure ϕ satisfying $\phi^4 \pm \phi^2 = 0$ Tensor, N.S., vol. 28, pp. 78-82.
- [3] K.Yano (1963).
 On a structure defined by a tensor field f of type (1, 1) satisfying f³ + f = 0.
 Tensor, N.S., vol. 14, pp. 99-109.
- K.Yano and M.Kon (1984).
 Structures on manifolds.
 World Scientific Publishing Company, Pte, Ltd., Farrer Road, Singapore, 9128.
- [5] K.Yano, S.Ishihara (1973)Tangent and Cotangent Bundles, Marcel Dekker Inc., New York,
- [6] F. Gouli-Andreou (1983) On Integrability conditions of a structure satisfying $f^5 + f = 0$, Tensor, Vol. 40(1), pp. 27-31.